This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.





Conversion of cysteine to S-sulfocysteine by reaction with sodium tetrathionate and reversal by exogenous thiols

FIG. 1A

The Cleavage of disulfide bonds by sodium sulfite to form the S-sulfo derivative

FIG. 1B



Preparation and Washing of TnI-containing Inclusion Bodies






FIG. 2




Summary of rTroponin-I Preparation – 3L5 (050900-051200)













SDS-PAGE Analysis Troponin Lot 3L5
Anion Exchange Steps #1 and #2
16% Tris-glycine Gel, Non-reducing, 5/15/00



- A. Sulfitolyzed Troponin Lot 3L4 Standard
- B. Solublized Inclusion Bodies
- C. Sulfitolyzed Inclusion Bodies (AEX #1 Load)
- D. Anion Exchange #1 Flowthrough
- E. Anion Exchange #1 Salt Eluate
- F. Anion Exchange #2 Load
- G. Anion Exchange #2 Flowthrough
- H. Anion Exchange #2 100 mM NaCl Eluate


FIG. 6







SDS-PAGE Analysis Troponin Lot 3L5 Hydrophobic Interaction Chromatography 16% Tris-glycine Gel, Non-reducing, 5/15/00



- A. Sulfitolyzed Troponin Lot 3L4 Standard
- B. AEX Step #2, Troponin Eluate Pool
- C. HIC Load, (w/1M Ammonium Sulfate)
- D. HIC Flowthrough (Troponin Product)
- E. HIC Low Salt Eluate (Column Strip)
- F. Lot 3L5 Sulfitolyzed Troponin Product

FIG. 8



Acq. Method : TROPC3Q.M

Analysis Method : C:\HPCHEM\1\METHODS\TROPC3Q.M
Last Changed : 03:04:24 pm

(modified after loading)

Quantitation of rTnI on Zorbax C3




FIG. 9



troponin I

LysC mapping

#	RC	Mass	Sequence
1	-2.1	707.28	- MGDEEK (1,6)
2	5.2	1618.96	- RNRAITARROHLK (7,19)
2 3	30.9	1431.76	- SVMLQIAATELEK (20,32)
4	-4.8	1132.55	- EESRREAEK (33,41)
5	48.0	2824.26	- QNYLAEHCPPLHIPGSMSEVQELCK (42,66)
6	-1.8	595.34	- QLHAK (67,71)
7	5.1	903.42	- IDAAEEEK (72,79)
5 6 7 8 9	10.9	1166.58	- YDMEVRVQK (80,88)
9	-9.4	334.19	- TSK (89,91)
10	5.6	1005.44	- ELEDMNQK (92,99)
11	13.9	847.49	- LFDLRGK (100, 106)
12	-1.7	293.17	- FK (107,108)
13	28.0	1896.07	
14	8.0	587.36	- ALLGSK (125,130)
15	-11.9	283.16	- HK (131,132)
16	20.5	1161.60	- VCMDLRANLK (133,142)
17	-4.8	373.23	- QVK (143,145)
18	-9.8	146.11	- K (146,146)
19	-6.8	620.27	- EDTEK (147,151)
20	15.7	1543.79	- ERDLRDVGDWRK (152,163)
21	-0.8	631.32	- NIEEK (164,168)
22	-4.4	763.36	- SGMEGRK (169,175)
23	-9.8	146.11	- K (176, 176)
24	7.7	728.27	- MFESES (177,182)

Current Chromatogram(s)

20




FIG. 10

40

45 min



SDS-PAGE Analysis Sulfitolyzed Troponin Reduction w/DTT 45 Minutes, Ambient Temperature, 1 mg.ml Tnl 6M Urea, 25 mM Tris, 0.15M NaCl pH 7.5 01/08/2000, 16% Tris-glycine Gel 2 3 4 5 6 7 8 9 10



1., 10., Mark 12 MW Stds 5. 0.2 mM DTT

2., 9., Sulfitolyzed Tn I 6. 0.3 mM DTT

3. 0.05 mM DTT

7. 0.5 mM DTT

4. 0.10 mM DTT

8. 1.0 mM DTT

FIG. 11